

$J/\psi\gamma$ and $\psi(2S)\gamma$ decay modes of the $X(3872)$

Yubing Dong^{1,2,3}, Amand Faessler³, Thomas Gutsche³, Valery E. Lyubovitskij^{3*}

¹ *Institute of High Energy Physics,
Beijing 100049, P. R. China*

² *Theoretical Physics Center for Science Facilities (TPCSF),
CAS, Beijing 100049, P. R. China*

³ *Institut für Theoretische Physik, Universität Tübingen,
Kepler Center for Astro and Particle Physics,
Auf der Morgenstelle 14, D-72076 Tübingen, Germany*

(Dated: September 2, 2009)

The $X(3872)$ with quantum numbers $J^{PC} = 1^{++}$ is considered as a composite state containing both molecular hadronic and a $c\bar{c}$ component. Based on this structure assumption we determine the radiative decay widths of the transitions $X(3872) \rightarrow J/\psi\gamma$ and $X(3872) \rightarrow \psi(2S)\gamma$. We find that in the $X(3872) \rightarrow J/\psi\gamma$ transition a destructive interference between the $c\bar{c}$ and hadronic components is needed to explain the large value of the ratio $\text{Br}(X(3872) \rightarrow \psi(2S)\gamma)/\text{Br}(X(3872) \rightarrow J/\psi\gamma)$ recently determined by the *BABAR* Collaboration.

PACS numbers: 12.38.Lg, 12.39.Fe, 13.25.Jx, 14.40.Gx

Keywords: charmonium, hadronic molecule, $X(3872)$, $\psi(2S)$, radiative decay

* On leave of absence from Department of Physics, Tomsk State University, 634050 Tomsk, Russia

I. INTRODUCTION

The $X(3872)$ is one of the peculiar new meson resonances which was discovered during the last years [1] and where its properties cannot be simply explained in the context of conventional constituent quark models. Presently several structure interpretations for this new resonance are proposed in the literature (for a status report see e.g. Refs. [2, 3, 4]). In the molecular approaches of [5]–[33] it is argued that the $X(3872)$ can be identified with a weakly-bound hadronic molecule whose constituents are D and D^* mesons. This natural interpretation is due to the fact that its mass m_X is very close to the $D^0\bar{D}^{*0}$ threshold and hence is in analogy to the deuteron — a weakly-bound state of proton and neutron.

First it was proposed that the state $X(3872)$ is a superposition of $D^0\bar{D}^{*0}$ and $\bar{D}^0D^{*0}$ pairs. Later (see e.g. discussions in Refs. [14, 16, 18]) other additional configurations such as a charmonium or even other meson pair components were discussed in addition to $D^0\bar{D}^{*0}$ + its charge conjugate (c.c.) (Here and in the following we use the convention that $\bar{D}^{*0}$ does not change sign under charge conjugation. For a detailed discussion see also Ref. [32]). The additional possibility of two nearly degenerate $X(3872)$ states with positive and negative charge parity has been discussed in Refs. [25, 34].

In the present paper we focus on the radiative decays of the $X(3872)$. There exist already several calculations for the radiative decays of the $X(3872)$ and various approaches give quite different results for the partial decay widths even when based on the same structure assumption. A new measurement by the *BABAR* Collaboration gives clear evidence for a strong radiative decay mode involving the $\psi(2S)$ [35]. They indicate the measured ratio of

$$\frac{\mathcal{B}(X(3872) \rightarrow \psi(2S)\gamma)}{\mathcal{B}(X(3872) \rightarrow J/\psi\gamma)} = 3.5 \pm 1.4. \quad (1)$$

As known from previous calculations in the molecular approach the radiative decay width of the $X(3872) \rightarrow \psi(2S)\gamma$ is always smaller than the one involving $J/\psi\gamma$. It is therefore expected that the ratio of Eq. (1) gives some constraint on a possible charmonium component in the $X(3872)$.

In Refs. [27, 31, 36, 37] we developed a formalism for the study of recently observed exotic meson states (like $D_{s0}^*(2317)$, $D_{s1}(2460)$, $X(3872)$, $\dots$) as hadronic molecules. In Ref. [31] we extended our formalism to the description of the decay $X \rightarrow J/\psi\gamma$ assuming that the X is a S -wave ($D^0\bar{D}^{*0} + D^{*0}\bar{D}^0$)/ $\sqrt{2}$ molecule of positive charge parity. As in the case of the D_{s0}^* and D_{s1} states a composite (molecular) structure of the $X(3872)$ meson is defined by the compositeness condition $Z = 0$ [38, 39, 40] (see also Refs. [27, 31, 36, 37]). This condition implies that the renormalization constant of the hadron wave function is set equal to zero or that the hadron exists as a bound state of its constituents. The compositeness condition was originally applied to the study of the deuteron as a bound state of proton and neutron [38]. Then it was extensively used in low-energy hadron phenomenology as the master equation for the treatment of mesons and baryons as bound states of light and heavy constituent quarks (see e.g. Refs. [39, 40]). By constructing a phenomenological Lagrangian including the couplings of the bound state to its constituents and of the constituents to other particles in the possible decay channels we calculated meson-loop diagrams describing different decays of the molecular states (see details in [27, 31, 36, 37]). In Ref. [31] we also estimated the role of a possible charmonium component in the $X(3872)$. We showed that the charmonium contribution to the $X \rightarrow J/\psi\gamma$ decay width is suppressed relative to the one of the molecular $D^0\bar{D}^{*0}$ component. In Ref. [27] we then considered the $X(3872)$ as a superposition of the molecular $D^0\bar{D}^{*0}$ component and other hadronic pairs — $D^\pm D^{*\mp}$, $J/\psi\omega$ and $J/\psi\rho$. Because the $D^0\bar{D}^{*0}$ component dominates in the transitions of X into charmonium states χ_{cJ} and pions we estimated these decays by restricting to this component only. In the analysis of decay modes involving J/ψ in the final state we used effective couplings for $XJ/\psi\omega$ and for $XJ/\psi\rho$ deduced from the compositeness condition. We also pointed out that an inclusion of a charmonium component, having destructive interference with the molecular components, can help to understand the ratio of $X \rightarrow \psi(2S)\gamma$ and $X \rightarrow J/\psi\gamma$ decays.

In the present paper we proceed as follows. In Sec. II we first discuss the basic notions of our approach. We discuss the effective Lagrangian for the treatment of the $X(3872)$ meson as a superposition of the molecular and charmonium components. Second, we consider the radiative two-body decays $X \rightarrow \psi(2S)\gamma$ and $X \rightarrow J/\psi\gamma$ decays. In Sec. III we present our numerical results. Finally, in Sec. IV we present a short summary.

II. APPROACH

Following Refs. [14, 27, 31] we consider the $X(3872)$ state as a superposition of the dominant molecular $D^0 D^{*0}$ component, of other hadronic configurations— $D^\pm D^{*\mp}$, $J/\psi\omega$, $J/\psi\rho$, as well as of the $c\bar{c}$ charmonium configuration as

$$|X(3872)\rangle = \cos\theta \left[\frac{Z_{D^0 D^{*0}}^{1/2}}{\sqrt{2}} (|D^0 \bar{D}^{*0}\rangle + |D^{*0} \bar{D}^0\rangle) + \frac{Z_{D^\pm D^{*\mp}}^{1/2}}{\sqrt{2}} (|D^+ D^{*-}\rangle + |D^- D^{*+}\rangle) + Z_{J\psi\omega}^{1/2} |J\psi\omega\rangle + Z_{J\psi\rho}^{1/2} |J\psi\rho\rangle \right] + \sin\theta c\bar{c}. \quad (2)$$

Here θ is the mixing angle between the hadronic and the charmonium components: $\cos^2\theta$ and $\sin^2\theta$ represent the probabilities to find a hadronic and charmonium configuration, respectively, for the normalization

$$Z_{D^0 D^{*0}} + Z_{D^\pm D^{*\mp}} + Z_{J\psi\omega} + Z_{J\psi\rho} = 1. \quad (3)$$

The limiting case of $\cos\theta = 1$ corresponds to the situation where $X(3872)$ has no charmonium component (this case was considered recently in Ref. [27]) while the value $\cos\theta = 0$ refers to the pure charmonium interpretation of the $X(3872)$. In this paper we will simply employ values for $Z_{D^0 D^{*0}}$, $Z_{D^\pm D^{*\mp}}$, $Z_{J\psi\omega}$ and $Z_{J\psi\rho}$ as derived in a potential model [14].

Our approach is based on an effective interaction Lagrangian describing the couplings of the $X(3872)$ to its constituents. Following our formalism developed in [27] we apply a nonlocal form containing the correlation functions $\Phi(y^2)$ characterizing the distribution of the constituents in the $X(3872)$. The Lagrangian is set up as:

$$\begin{aligned} \mathcal{L}_X(x) = & g_{X D^0 D^{*0}} X_\mu(x) J_{D^0 D^{*0}}^\mu(x) + g_{X D^\pm D^{*\mp}} X_\mu(x) J_{D^\pm D^{*\mp}}^\mu(x) \\ & + \frac{g_{X J\psi\omega}}{m_X} \epsilon_{\mu\nu\alpha\beta} \partial^\nu X^\alpha(x) J_{J\psi\omega}^{\mu\beta}(x) + \frac{g_{X J\psi\rho}}{m_X} \epsilon_{\mu\nu\alpha\beta} \partial^\nu X^\alpha(x) J_{J\psi\rho}^{\mu\beta}(x) \\ & + g_{X c\bar{c}} X_\mu(x) J_{c\bar{c}}^\mu(x), \end{aligned} \quad (4)$$

where $g_{X H_1 H_2}$ and $g_{X c\bar{c}}$ are the couplings of $X(3872)$ to the hadronic constituents $H_1 H_2$ and to the $c\bar{c}$ component. X is the field describing $X(3872)$ and $J_{\psi_1 \psi_2}^\Gamma$ are the respective currents composed of the fields ψ_1 and ψ_2 with:

$$J_{D\bar{D}^*}^\mu(x) = \frac{1}{\sqrt{2}} \int d^4 y \Phi_{D\bar{D}^*}(y^2) \left(D(x+y/2) \bar{D}^{*\mu}(x-y/2) + \bar{D}(x+y/2) D^{*\mu}(x-y/2) \right), \quad (5)$$

$$J_{J\psi V}^\mu(x) = \int d^4 y \Phi_V(y^2) V^\mu(x+y), \quad (6)$$

$$J_{c\bar{c}}^\mu(x) = \int d^4 y \Phi_{c\bar{c}}(y^2) \bar{c}(x+y/2) \gamma^\mu \gamma^5 c(x-y/2). \quad (7)$$

Here $\Phi_{D\bar{D}^*}$ is the correlation function describing the distribution of DD^* inside X , the function Φ_V describes the distribution of the light vector meson $V = \rho$ or ω around the J/ψ , which is located at the center of mass of the $X(3872)$, and the function $\Phi_{c\bar{c}}$ represents the distribution of the $c\bar{c}$ component. A basic requirement for the choice of an explicit form of the correlation functions Φ_I ($I = DD^*, V, c\bar{c}$) is that their Fourier transforms vanish sufficiently fast in the ultraviolet region of Euclidean space to render the Feynman diagrams ultraviolet finite. We adopt the same Gaussian form for all three correlation functions $\Phi_{DD^*} = \Phi_V = \Phi_{c\bar{c}} \equiv \Phi_X$ in order to reduce the number of free parameters. The Fourier transform of Φ_X is the universal vertex

$$\tilde{\Phi}_X(p_E^2/\Lambda_X^2) \doteq \exp(-p_E^2/\Lambda_X^2), \quad (8)$$

where p_E is the Euclidean Jacobi momentum and Λ_X is a size parameter. In the present paper we use the value of $\Lambda_X = 2$ GeV for all three correlation functions as fixed in Ref. [27].

The coupling constants $g_{X\psi_1\psi_2}$ are determined by the compositeness condition [36, 38, 39, 40]. This requirement implies that the renormalization constant of the hadron wave function is set equal to zero with

$$Z_X = 1 - \Sigma'_X(m_X^2) = 0. \quad (9)$$

Here $\Sigma'_X(m_X^2) = d\Sigma_X(p^2)/dp^2|_{p^2=m_X^2}$ is the derivative of the transverse part of the mass operator $\Sigma_X^{\mu\nu}$ conventionally split into the transverse Σ_X and longitudinal Σ_X^L parts as:

$$\Sigma_X^{\mu\nu}(p) = g_\perp^{\mu\nu} \Sigma_X(p^2) + \frac{p^\mu p^\nu}{p^2} \Sigma_X^L(p^2), \quad (10)$$

where $g_{\perp}^{\mu\nu} = g^{\mu\nu} - p^{\mu}p^{\nu}/p^2$ and $g_{\perp}^{\mu\nu}p_{\mu} = 0$. The mass operator of the $X(3872)$ receives contributions from five hadron-loop diagrams with

$$\Sigma_X(m_X^2) = \Sigma_{D^0 D^{*0}}(m_X^2) + \Sigma_{D^{\pm} D^{*\mp}}(m_X^2) + \Sigma_{J/\psi \omega}(m_X^2) + \Sigma_{J/\psi \rho}(m_X^2) + \Sigma_{c\bar{c}}(m_X^2), \quad (11)$$

which are induced by the interaction of X with the corresponding hadronic pairs $H_1 H_2$ and the $c\bar{c}$ configuration as contained in Eq. (4). The relevant diagram contributing to $\Sigma_X^{\mu\nu}(p)$ is shown in Fig.1. Using Eq. (2) and the compositeness condition (9) we have five independent equations to determine the coupling constants $g_{X\psi_1\psi_2}$.

In Fig.2 we display the diagrams relevant for the radiative decays $X \rightarrow J/\psi\gamma$ and $X \rightarrow \psi(2S)\gamma$. The main difference between the sets of diagrams feeding the channels $J/\psi\gamma$ and $\psi(2S)\gamma$ is: the last diagram of Fig.2 which is generated by the $J/\psi V$ ($V = \rho, \omega$) component of the $X(3872)$ only contributes to the $J/\psi\gamma$ final state. The effective Lagrangians generating the electromagnetic interaction vertices are

$$\begin{aligned} \mathcal{L}_{DD\gamma}(x) &= ieA_{\mu}(x) D^{-}(x) \overleftrightarrow{\partial}^{\mu} D^{+}(x), \\ \mathcal{L}_{D^{*}D^{*}\gamma}(x) &= ieA_{\mu}(x) \left\{ g^{\alpha\beta} D_{\alpha}^{*-} \overleftrightarrow{\partial}^{\mu} D_{\beta}^{*+}(x) + g^{\mu\beta} D_{\alpha}^{*-}(x) \partial^{\alpha} D_{\beta}^{*+}(x) - g^{\mu\alpha} \partial^{\beta} D_{\alpha}^{*-}(x) D_{\beta}^{*+}(x) \right\}, \\ \mathcal{L}_{D^{*}D\gamma}(x) &= \frac{e}{4} \epsilon^{\mu\nu\alpha\beta} F_{\mu\nu}(x) \left\{ g_{D^{*}-D+\gamma} D_{\alpha\beta}^{*-}(x) D^{+}(x) + g_{D^{*0}D^0\gamma} \bar{D}_{\alpha\beta}^{*0}(x) D^0(x) \right\} + \text{H.c.}, \\ \mathcal{L}_{cc\gamma} &= \frac{2e}{3} A_{\mu}(x) \bar{c}(x) \gamma^{\mu} c(x), \end{aligned} \quad (12)$$

where $A \overleftrightarrow{\partial}_{\mu} B = A \partial_{\mu} B - \partial_{\mu} A B$; $F_{\mu\nu} = \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu}$, and $M_{\mu\nu} = \partial_{\mu} M_{\nu} - \partial_{\nu} M_{\mu}$. The strong couplings of the charmonium states J/ψ and $\psi(2S)$ to D and D^{*} mesons are taken from heavy hadron chiral perturbation theory (HHChPT) [41] (see also Ref. [27]). For convenience we use a relativistic normalization of the meson states and write the Lagrangians in manifestly Lorentz covariant form:

$$\begin{aligned} \mathcal{L}_{\psi_n DD}(x) &= -ig_{\psi_n DD} \psi_n^{\mu} D_i^{\dagger}(x) \overleftrightarrow{\partial}_{\mu} D_i(x), \\ \mathcal{L}_{\psi_n DD^{*}}(x) &= g_{\psi_n DD^{*}} \epsilon_{\mu\nu\alpha\beta} \partial^{\mu} \psi_n^{\nu} \left\{ D_i^{*\beta\dagger}(x) \overleftrightarrow{\partial}^{\alpha} D_i(x) - D_i^{\dagger}(x) \overleftrightarrow{\partial}^{\alpha} D_i^{*\beta}(x) \right\}, \\ \mathcal{L}_{\psi_n D^{*}D^{*}}(x) &= ig_{\psi_n D^{*}D^{*}} \psi_n^{\mu} \left\{ -D_i^{*+\alpha} \overleftrightarrow{\partial}_{\mu} D_{i\alpha}^{*} + D_{i\mu}^{*+} \overleftrightarrow{\partial}_{\alpha} D_i^{*\alpha} + D_i^{*+\alpha} \overleftrightarrow{\partial}_{\alpha} D_{i\mu}^{*} \right\}, \end{aligned} \quad (13)$$

where ψ_n with $n = 1, 2$ stands for the vector fields of J/ψ and $\psi(2S)$, respectively. The chiral couplings of ψ_n with $D(D^{*})$ mesons read

$$g_{\psi_n DD} = g_{\psi_n D^{*}D^{*}} \frac{m_D}{m_{D^{*}}} = g_{\psi_n DD^{*}} m_{\psi_n} \sqrt{\frac{m_D}{m_{D^{*}}}} = \frac{m_{\psi_n}}{f_{\psi_n}}. \quad (14)$$

The quantity f_{ψ_n} is determined by the leptonic decay widths of J/ψ and $\psi(2S)$ of

$$\begin{aligned} \Gamma(J/\psi \rightarrow e^{+}e^{-}) &= \frac{16\pi}{27} \frac{\alpha^2}{m_{J/\psi}} f_{J/\psi}^2 = 5.55 \text{ keV}, \\ \Gamma(\psi(2S) \rightarrow e^{+}e^{-}) &= \frac{16\pi}{27} \frac{\alpha^2}{m_{\psi(2S)}} f_{\psi(2S)}^2 = 2.38 \text{ keV}, \end{aligned} \quad (15)$$

where α is the fine structure constant. Thus the ratios of the coupling constants $g_{\psi(2S)DD}$ and $g_{\psi(2S)D^{*}D^{*}}$ to $g_{J/\psi DD}$ and $g_{J/\psi D^{*}D^{*}}$ are fixed as

$$\frac{g_{\psi(2S)DD}}{g_{J/\psi DD}} = \frac{g_{\psi(2S)D^{*}D^{*}}}{g_{J/\psi D^{*}D^{*}}} = \frac{m_{\psi(2S)}}{m_{J/\psi}} \frac{f_{J/\psi}}{f_{\psi(2S)}} \simeq 1.67. \quad (16)$$

The coupling constant $g_{D^{*}D\gamma}$ is fixed by data (central values) on the radiative decay widths $\Gamma(D^{*} \rightarrow D\gamma)$ [1] with:

$$\begin{aligned} \Gamma(D^{*+} \rightarrow D^{+}\gamma) &= \frac{\alpha}{24} g_{D^{*}-D+\gamma}^2 m_{D^{*+}}^3 \left(1 - \frac{m_{D^{*+}}^2}{m_{D^{*+}}^2} \right)^3 = 1.54 \text{ keV}, \\ \Gamma(D^{*0} \rightarrow D^0\gamma) &= \frac{\alpha}{24} g_{D^{*0}D^0\gamma}^2 m_{D^{*0}}^3 \left(1 - \frac{m_{D^{*0}}^2}{m_{D^{*0}}^2} \right)^3 = 26.04 \text{ keV}. \end{aligned} \quad (17)$$

From Eq. (17) we finally predict

$$g_{D^{*-}D^+\gamma} = 0.5 \text{ GeV}^{-1}, \quad g_{D^{*0}D^0\gamma} = 2.0 \text{ GeV}^{-1}. \quad (18)$$

In our calculation the mass of the charm quark is chosen as $m_c \simeq M_X/2$. The respective couplings $g_{V\gamma}$ ($V = \rho^0, \omega$) of the transitions $\rho\gamma$ and $\omega\gamma$ are fixed from data on the $V \rightarrow e^+e^-$ decay widths as

$$\begin{aligned} \Gamma(\rho^0 \rightarrow e^+e^-) &= \frac{4\pi}{3} \alpha^2 g_{\rho\gamma}^2 m_\rho = 7.04 \text{ keV}, \\ \Gamma(\omega \rightarrow e^+e^-) &= \frac{4\pi}{3} \alpha^2 g_{\omega\gamma}^2 m_\omega = 0.60 \text{ keV}. \end{aligned} \quad (19)$$

Using last equation we get $g_{\rho\gamma} = 0.20$ and $g_{\omega\gamma} = 0.06$. Finally, the interaction Lagrangian for the coupling of ψ_n to the $c\bar{c}$ configuration is

$$\mathcal{L}_{\psi_n c\bar{c}}(x) = g_{\psi_n} \psi_n^\mu(x) \bar{c}(x) \gamma_\mu c(x), \quad (20)$$

where the coupling constant is given as

$$g_{\psi_n} = \frac{2}{3} \frac{m_{\psi_n}}{f_{\psi_n}}. \quad (21)$$

By using the decay constants f_{ψ_n} of the J/ψ and $\psi(2S)$ states we obtain the values $g_{J/\psi} = 4.96$ and $g_{\psi(2S)} = 8.26$.

III. RESULTS

To calculate the decay widths of the two processes $X(3872) \rightarrow J/\psi + \gamma$ and $X(3872) \rightarrow \psi(2S) + \gamma$ we further choose a typical value of 2 GeV for the free parameter Λ_X [27]. Here the binding energy $\epsilon = \epsilon_{D^0 D^{*0}}$ is defined by setting

$$M_X = M_{D^0} + M_{D^{*0}} - \epsilon_{D^0 D^{*0}}. \quad (22)$$

We present our numerical results for three typical values of the binding energy $\epsilon = 0.3, 0.7, 1$ MeV and use the corresponding sets of configuration probabilities as calculated in Ref. [14]:

$$Z_{D^0 D^{*0}} = 0.92, \quad Z_{D^\pm D^{*\mp}} = 0.033, \quad Z_{J_\psi \omega} = 0.041, \quad Z_{J_\psi \rho} = 0.006 \quad (23)$$

for $\epsilon = 0.3$ MeV,

$$Z_{D^0 D^{*0}} = 0.816, \quad Z_{D^\pm D^{*\mp}} = 0.079, \quad Z_{J_\psi \omega} = 0.096, \quad Z_{J_\psi \rho} = 0.009 \quad (24)$$

for $\epsilon = 0.7$ MeV, and

$$Z_{D^0 D^{*0}} = 0.758, \quad Z_{D^\pm D^{*\mp}} = 0.111, \quad Z_{J_\psi \omega} = 0.122, \quad Z_{J_\psi \rho} = 0.009 \quad (25)$$

for $\epsilon = 1$ MeV.

Fig. 3 contains the calculated results for the ratio

$$R_{2S} = \frac{\mathcal{B}(X(3872) \rightarrow \psi(2S) + \gamma)}{\mathcal{B}(X(3872) \rightarrow J/\psi + \gamma)} \quad (26)$$

in dependence on the mixing angle θ controlling the admixture of the $c\bar{c}$ component. For the binding energy ϵ we choose the discrete values indicated above. The experimental value including errors is reflected by the shadowed area. Moreover in Fig.4 we give predictions for the radiative decay widths of the two decay modes considered for the case of $\epsilon = 0.3$ MeV. Again we display the dependence of this quantities on the admixture of the $c\bar{c}$ component.

Since the charged D mesons are heavier than the neutral ones, the energy difference $\epsilon_{D^\pm D^{*\mp}} = M_{D^\pm} + M_{D^{*\mp}} - M_X$ is larger than the binding energy ϵ and we therefore get $g_{XD^0 D^{*0}} < g_{XD^\pm D^{*\mp}}$. Consequently the inclusion of the charged $D^\pm D^{*\mp}$ component leads to an increase in the radiative decay widths of the $X(3872)$. Our numerical calculation also shows that the presence of a $J/\psi V$ component in the $X(3872)$ is favored by the measured ratio R_{2S} . This can be traced to the fact that the decay mode $J/\psi\gamma$ (or $\psi(2S)\gamma$) is allowed (or forbidden) by the two components and they provide a subtract contribution to the widths of DD^* components. With respect to the presence of a possible $c\bar{c}$ component in the $X(3872)$ we note the following as based on our results. For the limiting cases of $\cos\theta = 1$ or 0 one

cannot reasonably explain the large ratio of R_{2S} . This is mainly due to the fact that the phase space for the $J/\psi\gamma$ transition (with three-momentum of the outgoing photon $\mathbf{p} = 654$ MeV) is much larger than that of $\psi(2S)\gamma$ (with the photon momentum $\mathbf{p} = 182$ MeV). However the results of Fig.3 indicate that a small admixture of a $c\bar{c}$ component is essential to reproduce the large measured value for R_{2S} . This feature is due to the destructive interference between the $J/\psi\gamma$ decay amplitudes arising from the $c\bar{c}$ and the hadronic molecular components. Quantitatively this effect is also visible in the prediction for the decay width $X \rightarrow J/\psi\gamma$ of Fig.4. Also, as we already showed in Ref. [27] the inclusion of the $J/\psi V$ components in the structure of the $X(3872)$ leads to a decrease in the $X \rightarrow J/\psi\gamma$ rate because $\psi(2S)V$ components are absent in the $X(3872)$. From Fig.3 we deduce that the experimental value of R_{2S} is reproduced for a small admixture of the $c\bar{c}$ component with $\sin\theta \simeq -(0.16 - 0.24)$. An enlarged value for the binding energy ϵ also leads to an increase in the $c\bar{c}$ probability requiring that the data are reproduced. If no $c\bar{c}$ component is present in the $X(3872)$ the decay width for $J/\psi\gamma$ is about 60 keV and is much larger than the one for the $\psi(2S)\gamma$ channel (about 0.3 keV). The addition of a $c\bar{c}$ admixture leads to a monotonous increase in the $\psi(2S)\gamma$ decay width. At the same time the decay width for $J/\psi\gamma$ decreases rapidly in the region of $\sin\theta \geq -0.16$ due to this destructive interference and then it increases further. Therefore, for mixing values of about $\sin\theta \geq -0.16$ in the case of $\epsilon = 0.3 - 1$ MeV the decay width for $\psi(2S)\gamma$ may exceed the one of $J/\psi\gamma$ and a consistent explanation for the observed ratio R_{2S} is obtained. In this case the radiative decay widths are about 1 – 2 keV for $J/\psi\gamma$ and 5 – 6 keV for $\psi(2S)\gamma$, respectively.

IV. SUMMARY

In summary, we have shown that inclusion of both the hadronic component $J/\psi V$ and the $c\bar{c}$ charmonium configuration in the $X(3872)$ structure is necessary to reproduce the large value of the ratio R_{2S} of the $X(3872)$ measured by the *BABAR* Collaboration. In our calculation we only need a small admixture of the $c\bar{c}$ component with $\sin\theta \simeq -(0.16 - 0.24)$ and the binding energy ϵ ranging from 0.3 to 1 MeV to explain the ratio R_{2S} . Moreover we found that after $c\bar{c}$ admixture the radiative decay width of $J/\psi\gamma$ decreases to about 1 – 2 keV while the measured ratio of R_{2S} is reasonably explained. We therefore conclude that both the measured value of the ratio R_{2S} and the yet to be experimentally determined value for $\Gamma(X(3872) \rightarrow J/\psi\gamma)$ provide a constraint on the $c\bar{c}$ component in the $X(3872)$.

Acknowledgments

This work was supported by the DFG under Contract No. FA67/31-2 and No. GRK683. This work is supported by the National Sciences Foundations No. 10775148 and by CAS grant No. KJCX3-SYW-N2 (YBD). This research is also part of the European Community-Research Infrastructure Integrating Activity “Study of Strongly Interacting Matter” (HadronPhysics2, Grant Agreement No. 227431) and of the President grant of Russia “Scientific Schools” No. 871.2008.2. The work is partially supported by Russian Science and Innovations Federal Agency under contract No 02.740.11.0238. Y.B.D. would like to thank the Tübingen theory group for its hospitality. V.E.L. would like to thank the Theory Group of the Institute of High Energy Physics (Beijing) for its hospitality.

-
- [1] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008).
 - [2] E. S. Swanson, Phys. Rept. **429**, 243 (2006) [arXiv:hep-ph/0601110].
 - [3] G. Bauer, Int. J. Mod. Phys. A **21**, 959 (2006) [arXiv:hep-ex/0505083].
 - [4] M. B. Voloshin, Prog. Part. Nucl. Phys. **61**, 455 (2008) [arXiv:0711.4556 [hep-ph]].
 - [5] M. B. Voloshin and L. B. Okun, JETP Lett. **23**, 333 (1976) [Pisma Zh. Eksp. Teor. Fiz. **23**, 369 (1976)].
 - [6] A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. Lett. **38**, 317 (1977).
 - [7] N. A. Tornqvist, Z. Phys. C **61**, 525 (1994) [arXiv:hep-ph/9310247].
 - [8] N. A. Tornqvist, Phys. Lett. B **590**, 209 (2004) [arXiv:hep-ph/0402237].
 - [9] S. Pakvasa and M. Suzuki, Phys. Lett. B **579**, 67 (2004) [arXiv:hep-ph/0309294].
 - [10] M. B. Voloshin, Phys. Lett. B **579**, 316 (2004) [arXiv:hep-ph/0309307].
 - [11] E. Braaten and M. Kusunoki, Phys. Rev. D **69**, 074005 (2004) [arXiv:hep-ph/0311147].
 - [12] F. E. Close and P. R. Page, Phys. Lett. B **578**, 119 (2004) [arXiv:hep-ph/0309253].
 - [13] T. Barnes and S. Godfrey, Phys. Rev. D **69**, 054008 (2004) [arXiv:hep-ph/0311162].
 - [14] E. S. Swanson, Phys. Lett. B **588**, 189 (2004) [arXiv:hep-ph/0311229].
 - [15] E. S. Swanson, Phys. Lett. B **598**, 197 (2004) [arXiv:hep-ph/0406080].

- [16] M. B. Voloshin, Phys. Lett. B **604**, 69 (2004) [arXiv:hep-ph/0408321].
- [17] E. Braaten and M. Kusunoki, Phys. Rev. D **72**, 014012 (2005) [arXiv:hep-ph/0506087].
- [18] E. Braaten and M. Kusunoki, Phys. Rev. D **72**, 054022 (2005) [arXiv:hep-ph/0507163].
- [19] M. T. AlFiky, F. Gabbiani and A. A. Petrov, Phys. Lett. B **640**, 238 (2006) [arXiv:hep-ph/0506141].
- [20] Yu. S. Kalashnikova, Phys. Rev. D **72**, 034010 (2005) [arXiv:hep-ph/0506270].
- [21] M. B. Voloshin, Int. J. Mod. Phys. A **21**, 1239 (2006) [arXiv:hep-ph/0509192].
- [22] E. Braaten and M. Lu, Phys. Rev. D **74**, 054020 (2006) [arXiv:hep-ph/0606115].
- [23] X. Liu, B. Zhang and S. L. Zhu, Phys. Lett. B **645**, 185 (2007) [arXiv:hep-ph/0610278]; C. Meng and K. T. Chao, Phys. Rev. D **75**, 114002 (2007) [arXiv:hep-ph/0703205].
- [24] E. Braaten, M. Lu and J. Lee, Phys. Rev. D **76**, 054010 (2007) [arXiv:hep-ph/0702128].
- [25] D. Gamermann and E. Oset, Eur. Phys. J. A **33**, 119 (2007) [arXiv:0704.2314 [hep-ph]].
- [26] D. Gamermann and E. Oset, Phys. Rev. D **80**, 014003 (2009) [arXiv:0905.0402 [hep-ph]].
- [27] Y. B. Dong, Amand Faessler, Thomas Gutsche, Sergey Kovalenko and Valery E. Lyubovitskij, Phys. Rev. D **79**, 094013 (2009) [arXiv:0903.5416 [hep-ph]].
- [28] Y. R. Liu, X. Liu, W. Z. Deng and S. L. Zhu, Eur. Phys. J. C **56**, 63 (2008) [arXiv:0801.3540 [hep-ph]].
- [29] S. Fleming, M. Kusunoki, T. Mehen and U. van Kolck, Phys. Rev. D **76**, 034006 (2007) [arXiv:hep-ph/0703168].
- [30] E. Braaten and M. Lu, Phys. Rev. D **77**, 014029 (2008) [arXiv:0710.5482 [hep-ph]].
- [31] Y. B. Dong, A. Faessler, T. Gutsche and V. E. Lyubovitskij, Phys. Rev. D **77**, 094013 (2008) [arXiv:0802.3610 [hep-ph]].
- [32] C. E. Thomas and F. E. Close, Phys. Rev. D **78**, 034007 (2008) [arXiv:0805.3653 [hep-ph]].
- [33] S. Fleming and T. Mehen, Phys. Rev. D **78**, 094019 (2008) [arXiv:0807.2674 [hep-ph]].
- [34] K. Terasaki, Prog. Theor. Phys. **118**, 821 (2007) [arXiv:0706.3944 [hep-ph]].
- [35] B. Aubert *et al.* (BABAR Collaboration), arXiv:0809.0042 [hep-ex].
- [36] A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, Phys. Rev. D **76**, 014005 (2007) [arXiv:0705.0254 [hep-ph]]; A. Faessler, T. Gutsche, S. Kovalenko and V. E. Lyubovitskij, Phys. Rev. D **76**, 014003 (2007) [arXiv:0705.0892 [hep-ph]]; A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, Phys. Rev. D **76**, 114008 (2007) [arXiv:0709.3946 [hep-ph]]; A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, Phys. Rev. D **77**, 114013 (2008) [arXiv:0801.2232 [hep-ph]].
- [37] T. Branz, T. Gutsche and V. E. Lyubovitskij, Phys. Rev. D **79**, 014035 (2009) [arXiv:0812.0942 [hep-ph]]; T. Branz, T. Gutsche and V. Lyubovitskij, arXiv:0905.3979 [hep-ph] (accepted for publication in Phys. Rev. D).
- [38] S. Weinberg, Phys. Rev. **130**, 776 (1963); A. Salam, Nuovo Cim. **25**, 224 (1962); K. Hayashi, M. Hirayama, T. Muta, N. Seto and T. Shirafuji, Fortsch. Phys. **15**, 625 (1967).
- [39] G. V. Efimov and M. A. Ivanov, *The Quark Confinement Model of Hadrons*, (IOP Publishing, Bristol & Philadelphia, 1993).
- [40] I. V. Anikin, M. A. Ivanov, N. B. Kulimanova and V. E. Lyubovitskij, Z. Phys. C **65**, 681 (1995); M. A. Ivanov, M. P. Locher and V. E. Lyubovitskij, Few Body Syst. **21**, 131 (1996); M. A. Ivanov, V. E. Lyubovitskij, J. G. Körner and P. Kroll, Phys. Rev. D **56**, 348 (1997) [arXiv:hep-ph/9612463]; M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij and A. G. Rusetsky, Phys. Rev. D **60**, 094002 (1999) [arXiv:hep-ph/9904421]; A. Faessler, T. Gutsche, M. A. Ivanov, V. E. Lyubovitskij and P. Wang, Phys. Rev. D **68**, 014011 (2003) [arXiv:hep-ph/0304031]; A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij, D. Nicmorus and K. Pumsa-ard, Phys. Rev. D **73**, 094013 (2006) [arXiv:hep-ph/0602193]; A. Faessler, T. Gutsche, B. R. Holstein, V. E. Lyubovitskij, D. Nicmorus and K. Pumsa-ard, Phys. Rev. D **74**, 074010 (2006) [arXiv:hep-ph/0608015]; A. Faessler, T. Gutsche, B. R. Holstein, M. A. Ivanov, J. G. Körner and V. E. Lyubovitskij, Phys. Rev. D **78**, 094005 (2008) [arXiv:0809.4159 [hep-ph]]; A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Körner and V. E. Lyubovitskij, Phys. Rev. D **80**, 034025 (2009) [arXiv:0907.0563 [hep-ph]].
- [41] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Rev. D **69**, 054023 (2004) [arXiv:hep-ph/0310084].




FIG. 1: $H_1 H_2$ hadron-loop or $c\bar{c}$ quark-loop diagram contributing to the mass operator of the $X(3872)$ meson.




FIG. 2: Diagrams contributing to the radiative transitions $X(3872) \rightarrow J/\psi + \gamma$ and $X(3872) \rightarrow \psi(2S) + \gamma$.




FIG. 3: Ratio R_{2S} . Solid, dashed and dotted-dashed lines are the results for $\epsilon = 0.3$ MeV, $\epsilon = 0.7$ MeV and $\epsilon = 1$ MeV, respectively.




FIG. 4: Radiative decay widths of the $X(3872)$ for $\epsilon = 0.3$ MeV. Solid and dotted curves are the results for the transitions $X \rightarrow J/\psi\gamma$ and $X \rightarrow \psi(2S)\gamma$, respectively.